



**QUEEN'S  
UNIVERSITY  
BELFAST**

## Climatic controls on Later Stone Age human adaptation in Africa's southern Cape

Chase, B. M., Faith, J. T., Mackay, A., Chevalier, M., Carr, A. S., Boom, A., ... Reimer, P. J. (2018). Climatic controls on Later Stone Age human adaptation in Africa's southern Cape. *Journal of Human Evolution*, 114, 35-44. <https://doi.org/10.1016/j.jhevol.2017.09.006>

### Published in:

Journal of Human Evolution

### Document Version:

Peer reviewed version

### Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

### Publisher rights

Copyright 2017 Elsevier.

This manuscript is distributed under a Creative Commons Attribution-NonCommercial-NoDerivs License

(<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

### General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

# Climatic controls on Later Stone Age human adaptation in Africa's southern Cape

---

Brian M. Chase<sup>\*1</sup>, J. Tyler Faith<sup>2</sup>, Alex Mackay<sup>3</sup>, Manuel Chevalier<sup>1</sup>, Andrew S. Carr<sup>4</sup>,  
Arnoud Boom<sup>4</sup>, Sophak Lim<sup>1</sup>, Paula J. Reimer<sup>5</sup>

<sup>1</sup>Centre National de la Recherche Scientifique, UMR 5554, Institut des Sciences de l'Evolution de Montpellier, Université Montpellier, Bat.22, CC061, Place Eugène Bataillon, 34095 Montpellier, cedex5, France;

<sup>2</sup>Archaeology Program, School of Social Science, University of Queensland, Brisbane, QLD 4072, Australia;

<sup>3</sup>Centre for Archaeological Science, School of Earth and Environmental Sciences, Northfields Avenue, Building 41, University of Wollongong, NSW 2522, Australia;

<sup>4</sup>Department of Geography, University of Leicester, Leicester, LE1 7RH, UK;

<sup>5</sup>School of Natural and Built Environment, Queen's University Belfast, Belfast, BT7 1NN, Northern Ireland, UK

\*Correspondence to:

Email: [brian.chase@univ-montp.fr](mailto:brian.chase@univ-montp.fr)

## Abstract

Africa's southern Cape is a key region for the evolution of our species, with early symbolic systems, marine faunal exploitation, and episodic production of microlithic stone tools taken as evidence for the appearance of distinctively complex human behavior. However, the temporally discontinuous nature of this evidence precludes ready assumptions of intrinsic adaptive benefit, and has encouraged diverse explanations for the occurrence of these behaviors, in terms of regional demographic, social and ecological conditions. Here, we present a new high-resolution multi-proxy record of environmental change that indicates that faunal exploitation patterns and lithic technologies track climatic variation across the last 22,300 years in the southern Cape. Conditions during the Last Glacial Maximum and deglaciation were humid, and zooarchaeological data indicate high foraging returns. By contrast, the Holocene is characterized by much drier conditions and a degraded resource base. Critically, we demonstrate that systems for technological delivery – or provisioning – were responsive to changing humidity and environmental productivity. However, in contrast to prevailing models, bladelet-rich microlithic technologies were deployed under conditions of high foraging returns and abandoned in response to increased aridity and less productive subsistence environments. This suggests that posited links between microlithic technologies and subsistence risk are not universal, and the behavioral sophistication of human populations is reflected in their adaptive flexibility rather than in the use of specific technological systems.

**Keywords:** paleoclimate; paleoecology; rock hyrax middens; microlithic; macrofauna; Boomplaas Cave

## Introduction

South Africa's southern coastal margin is a key region for the evolution and development of our species (Ambrose, 2002; Ambrose and Lorenz, 1990; Brown et al., 2012; Henshilwood et al., 2004a; Henshilwood et al., 2002; Marean, 2010; Powell et al., 2009). The southern Cape archaeological record has reframed the debate about the evolution of human behavior, providing early examples of engravings, ornaments, heat treatment of tool-stone and the focussed consumption of marine resources (Delagnes et al., 2016; Henshilwood et al., 2004b; Henshilwood et al., 2002; Henshilwood et al., 2014; Marean, 2014). The region also exhibits regular technological turnover through the last 100,000 years, with the intermittent production of bladelets, bifacial points and backed artefacts and the use of fine-grained rock, interspersed with periods lacking regular retouched flake forms and dominated by locally available rocks such as quartzite and quartz (Deacon, 1984; Wilkins et al., 2017). The links between these variable technological and subsistence records and their environmental context – necessary to arguments about the evolution of human adaptation – remain surprisingly unclear (Deacon, 1982; Roberts et al., 2016). This reflects the region's particular climatic dynamism (Chase and Meadows, 2007) coupled with disagreement concerning the interpretation of its paleoenvironmental archives (e.g. Chase and Meadows, 2007; Deacon and Lancaster, 1988; Faith, 2013b; Marean et al., 2014).

In this paper, we focus on the Later Stone Age record in the southern Cape, for which - in contrast with the Middle Stone Age - high resolution environmental and archaeological data are now available. We explore the strength of coupling between environments, subsistence behavior and lithic technology over the last 22,300 years to understand whether, and how closely, human behavior tracked environmental change. Spanning the transition from the Last Glacial Maximum (LGM; 26.5-19 ka; Clark et al., 2009) to the Holocene (11.7 ka to present Lowe et al., 2008), and episodes of the use of bladelet-rich technological systems, our data also have a bearing on broader debates about the role of what are often termed 'microlithic' technologies in issues of human adaptation and expansion.

## Later Stone Age environments and archaeology in the southern Cape

Influenced by both temperate and tropical climate systems (Figure 1), long-term climate change in the southern Cape is characterized by significant and often abrupt fluctuations (Bard and Rickaby, 2009; Chase et al., 2013; Chase and Meadows, 2007; Heaton et al., 1986;

Quick et al., 2015; Quick et al., 2016; Talma and Vogel, 1992). Existing evidence indicates that during the Holocene the relative influences of the two dominant synoptic scale moisture-bearing systems – 1) the southern westerly storm track, which expands/shifts northward in the winter, and 2) the tropical easterly flow, which transports moisture from the Indian Ocean during the summer – have varied significantly (Chase et al., 2013; Chase et al., 2015b). However, there is little detailed paleoenvironmental evidence pre-dating the Holocene (Carr et al., 2016b; Chase and Meadows, 2007), and as a result there are contradictory opinions concerning conditions since the LGM (Chase and Meadows, 2007; Deacon and Lancaster, 1988; Faith, 2013b; Kohfeld et al., 2013; Partridge et al., 1999; Partridge et al., 2004; Sime et al., 2013; Stone, 2014), to the extent that some studies conclude that the region was exceptionally “harsh” and arid during the LGM (Deacon and Lancaster, 1988; Scholtz, 1986), while others infer greater humidity and highly productive terrestrial environments (e.g. Faith, 2013b; Parkington et al., 2000). This uncertainty has fundamentally hindered our understanding of past climate dynamics in the region, and, by extension, the impact of past climate change on hunter-gatherer adaptive and subsistence strategies during both the Later and Middle Stone Age.

In the southern Cape, the Later Stone Age archaeological sequence is typically divided into several industries or technocomplexes: early Later Stone Age (ELSA ~<40-24 cal kBP), Robberg (~24-12 cal kBP), Oakhurst (~12-8 cal kBP) and Wilton (~8-2 cal kBP), followed by the arrival of Khoikhoi herders in the last 2000 years (Deacon et al., 1984; Deacon, 1978; Lombard et al., 2012; Mitchell, 1988). The ELSA is associated with the production of small flakes, often through bipolar reduction of cores, though it otherwise lacks unifying characteristics and has been described as a period of technological heterogeneity (Mitchell, 1988; Wadley, 1993). The Robberg presents more coherent characteristics, including the production of large numbers of bladelets (small, elongate flakes usually less than 24 mm long) produced both from dedicated bladelet cores and from those worked by bipolar reduction (Mitchell, 1988). The Robberg also sees more concentrated, if episodic, use of fine-grained rocks such as a silcrete and chert than the preceding or subsequent phases (Deacon, 1978; Deacon, 1982). The Oakhurst (or Albany) is typified by fewer bladelets, larger flakes, a range of scraper forms and declining use of fine-grained rock, while the Wilton features both scrapers and backed artefacts and highly variable patterns of raw material use (Deacon, 1972; Deacon, 1978; Lombard et al., 2012). While these units are

coarse and mask considerable variation, they provide a useful heuristic for discussing broad patterns in technological change across the later LSA.

Consistent with the imprecise meaning of the term (Pargeter, 2016), the ELSA, Robberg and Wilton have all been described as ‘microlithic’ (Bousman, 2005; Deacon, 1984; Mitchell, 1988; Wadley, 1993), but based on different characteristics – small flakes in the case of the ELSA, bladelets in the case of the Robberg and backed artefacts in the Wilton (Lombard et al., 2012). The advent of dedicated bladelet production in particular – as characterizes the Robberg – is argued to have presented humans with a significant adaptive advantage during our evolution and dispersal (Ambrose, 2002; Bar-Yosef and Kuhn, 1999; Clarkson et al., 2009; Foley and Lahr, 2003). Some researchers have linked an emphasis on bladelet production with responses to heightened subsistence risk associated with low or declining subsistence resource productivity (Elston and Brantingham, 2002; Petraglia et al., 2009) (for discussion of the risk concept used here see Bamforth and Bleed, 1997). Others have suggested that bladelet production provided benefits under conditions of high residential mobility (Goebel, 2002; Neeley, 2002). Both explanations – increased subsistence risk and increased mobility – have been posited for bladelet-rich systems in southern Africa during globally cooler conditions (Ambrose, 2002; Grosjean et al., 2003; McCall, 2007; McCall and Thomas, 2012; Mellars, 2006; Mitchell, 2000). The Robberg specifically has been associated with increased residential mobility in response to inferred diminishing resource density (Ambrose, 2002; Mitchell, 2000), and has been explained as a risk-dampening response to resource stress (Mackay, 2009). Other researchers, however, have suggested that any tracking between LSA technological systems and paleoenvironmental variation was relatively weak, and occurred only at the broadest scale of environmental change (e.g. Deacon, 1982). The reality of coupling between technology, subsistence conditions and environmental change in this period is thus contested, and with it the viability of high-order explanations for the behavioral significance of artefacts such as bladelets.

## **Sites and regional setting**

To explore the relationship between environmental change and human activities and technology, we focus on sites from the Swartberg mountains of South Africa’s southern Cape, one of the major ranges in the east-west axis of the Cape Fold Mountains (Figure 1). From Seweweekspoort, a deep transversal valley in the central Swartberg mountains, a series of rock hyrax (*Procavia capensis*) middens – stratified accumulations of dried urine and fecal

pellets (see Chase et al., 2012) – were identified, and six middens from two sites, Seweweekspoort-1 (SWP-1; 33.3668°S, 21.4144°E) and Seweweekspoort-3 (SWP-3; 33.4092°S, 21.4031°), were analysed for this study. SWP-1 is located on a west-facing cliff on the northern slope of the pass (Figure 1). The SWP-1 middens were taken from several locations within the same larger shelter, formed by a ~100 m overhanging cliff. SWP-3 is located on a low eastern cliff in the central section of the valley near the valley bottom, and experiences a more humid microclimate relative to the exposed position of SWP-1.

<insert Figure 1>

The SWP-1 site is located in the North Swartberg Sandstone Fynbos, but less than a kilometre to the north is the Matjiesfontein Shale Renosterveld (Mucina and Rutherford, 2006). The former, depending on altitude and aspect, is predominantly asteraceous, proteoid and restioid fynbos, while the latter is dominated by asteraceous elements, particularly *Elytropappus rhinocerotis*, *Eriocephalus* sp. and *Euryops* sp., and by an increasing number of succulents, primarily from the Crassulaceae family (Mucina and Rutherford, 2006). At SWP-1, these vegetation types inter-digitate to some extent, with the broad west-facing drainage just north of the site supporting more hydrophilic taxa, such as *Protea*. The north-facing rock shelves directly adjacent to the site maintain only shallow soils and a dominance of succulents. Grasses in the region are a mixture of C<sub>3</sub> (e.g., *Erharta*) and C<sub>4</sub> (e.g., *Aristida*, *Stipagrostis*) varieties (<http://sibis.sanbi.org/faces/DataSources.jsp>; Rutherford et al., 2012; Rutherford et al., 2003; SANBI, 2003). SWP-3 is located in the South Swartberg Sandstone Fynbos, which at the site is primarily proteoid in character. In addition, there are numerous arboreal/Cape thicket elements (e.g., *Podocarpus*, Celastraceae, *Dodonaea*, *Searsia*, *Euclea*, Oleaceae) associated with drainages and the nearby riparian zones of the valley bottom.

The Seweweekspoort sites are located 70 km west of the well-stratified late Pleistocene / Holocene archaeological site of Boomplaas Cave. This site is also situated on the flanks of the Swartberg and occupies a very similar climatic regime, making it ideal for exploring linkages between climate, environment, subsistence and technology at high temporal resolution. Located on the southern flanks of the Swartberg range overlooking the Cango Valley, Boomplaas Cave was excavated by Hilary Deacon in the 1970s (Deacon,

1979). The excavated sediments were divided into a series of members, some of which were subdivided into units. The site is positioned within a transitional shrubland whose component species vary as a function of temperature and moisture gradients moving upslope from the floor of the valley (Vlok and Schutte-Vlok, 2010). The lowlands to the south are characterized by renosterveld habitats, which give way to fynbos vegetation along the slopes of the Swartberg. Along watercourses and ravines in the Cango Valley are more densely wooded habitats that include sweet thorn trees (*Vachellia (Acacia) karroo*) and ironwood (*Olea* spp.) among others (Moffett and Deacon, 1977). As is also the case at Seweweekspoort, Boomplaas today receives contributions of both winter and summer rainfall (Figure 1). Thus, the limited grasses that do occur in the area today include a mixture of both C<sub>3</sub> and C<sub>4</sub> species.

## Materials and methods

Rock hyrax middens accumulate over thousands of years and preserve continuous records of past climate change (Chase et al., 2012). The six middens from Seweweekspoort sites SWP-1 and SWP-3 were selected for analysis because they are composed almost entirely of hyraceum (no visible fecal pellets). Our experience suggests that such middens have superior stratigraphic integrity compared to more pellet-rich middens. Representative portions of the middens were processed as described in Chase et al. (2013; 2012). Radiocarbon age determinations ( $n=36$ ) were processed at the <sup>14</sup>CHRONO Centre, Queen's University Belfast using accelerator mass spectrometry (AMS) (Supplementary Online Material [SOM] Figure S1; Table S1). The radiocarbon ages were corrected for isotope fractionation using the AMS measured  $\delta^{13}\text{C}$  and calibrated using the SHCal13 calibration data (Hogg et al., 2013). The Bacon 3.0.3 software package (Blaauw and Christen, 2011) was used to generate all age-depth models (SOM Figure S1). Results indicate that these sequences continuously span the last 22,300 years.

The fossil pollen content of 82 adjacent, contiguous pollen samples were prepared with standard physical (600  $\mu\text{m}$  sieving and decanting) and chemical (HCl, KOH, HF and acetolysis) methods (Moore et al., 1991). *Lycopodium* tablets were added to the weighed sample to estimate pollen concentrations (Stockmarr, 1971). A minimum pollen sum of 400 grains was counted at a magnification of  $\times 400$  under a light microscope, and identified with the help of the literature (Scott, 1982; van Zinderen Bakker, 1953, 1956; van Zinderen



Bakker and Coetzee, 1959), and photographic and slide reference collections at the Universities of the Free State, Cape Town, and Montpellier.

The bulk stable nitrogen ( $^{15}\text{N}$ ) and carbon ( $^{13}\text{C}$ ) isotope contents of 767 overlapping hyraceum samples were measured at the Department of Archaeology, University of Cape Town following Chase et al. (2010; 2009; 2011; 2012), with contiguous/overlapping samples obtained from two series of offset 1 mm holes. For the stable isotope analyses, the standard deviation derived from replicate analyses of homogeneous material was better than 0.2‰ for both nitrogen and carbon. Nitrogen isotope results are expressed relative to atmospheric nitrogen (SOM Figure S2). Carbon isotope results are expressed relative to Vienna PDB (SOM Figure S3).

Stable isotope results from the different Seweweekspoort rock hyrax middens were combined into a single aggregate record using Local Regression (LOESS) curve fitting of the combined datasets (SOM Figures S2 and S3). As individual middens under the same climate regime may exhibit differences in their isotopic records due to microclimatic influences on individual foraging ranges (i.e., baseline  $\delta^{15}\text{N}$  variability), we have adjusted the  $\delta^{15}\text{N}$  to account for these differences prior to LOESS curve fitting. Using the SWP-1-1 and SWP-1-4b records as a datum, an estimated offset of 1.5‰ was added to the  $\delta^{15}\text{N}$  data from the SWP-3-1 to compensate for the more humid microclimate in which the midden was found, and 0.5‰ and 1‰ were added to SWP-1-5 and SWP-1-2a respectively to account for their more exposed positions.

The carbon isotopic composition of the hyraceum is representative of vegetation around a midden site (Carr et al., 2016a) and provides information on 1) the relative contribution of  $\text{C}_3$ ,  $\text{C}_4$  and CAM plants (Smith, 1972) to the animals' diet, and 2) variations in plant water-use efficiency (WUE) as a function of climate (Ehleringer and Cooper, 1988; Farquhar et al., 1989; Farquhar and Richards, 1984; Pate, 2001). Throughout the broader region, the distribution of  $\text{C}_3$  and  $\text{C}_4$  grasses tracks the proportion of winter versus summer rainfall (Vogel, 1978). As mentioned, at Seweweekspoort today, grasses are a mosaic of  $\text{C}_3$  and  $\text{C}_4$  varieties (Rutherford et al., 2012; Rutherford et al., 2003; SANBI, 2003), and where aspect and soil depth limit soil water content, CAM plants become increasingly abundant. As  $\text{C}_3$  plants are depleted in  $^{13}\text{C}$  compared with most CAM and all  $\text{C}_4$  plants, higher  $\delta^{13}\text{C}$  values indicate more abundant warm season ( $\text{C}_4$ ) grasses and/or succulent plants (CAM), and generally warmer/more arid conditions.

Hyraceum  $\delta^{15}\text{N}$  is an indicator of changes in ecosystem water-availability (Carr et al., 2016a; Chase et al., 2013; Chase et al., 2015b; Chase et al., 2009; Chase et al., 2011). A positive relationship exists between aridity and  $\delta^{15}\text{N}$  in soils, plants and herbivores, with drier conditions correlating with enriched  $\delta^{15}\text{N}$  (Carr et al., 2016a), most likely as a result of denitrification processes in arid/semi-arid soils (Handley et al., 1999; Handley et al., 1994; Hartman, 2011; Heaton, 1987; Murphy and Bowman, 2006, 2009; Wang et al., 2010). In the hyraceum samples, the narrowly defined feeding range of the hyraxes (<60 m; Sale, 1965), and the accumulation rates of the middens (~20-60 years/sample) enforce a spatio-temporal averaging that reduces the  $\delta^{15}\text{N}$  variability observed in modern ecosystem studies (Carr et al., 2016a), and provides a more reliable index of past water variability (Carr et al., 2016a; Chase et al., 2012).

#### *Boomplaas faunal and archaeological archives*

The Boomplaas sequence spans much of the last >65,000 years (Deacon, 1982), though we focus here on the fauna and flaked stone artifacts from the upper stratigraphic units corresponding in age with the Seweweekspoort record (SOM Table S2). We use these data to explore the relationship between the paleoenvironmental changes documented at Seweweekspoort and mammal community composition, foraging efficiency, and technological organization. Ages for Boomplaas follow Deacon (1982), calibrated using SHCal13 (Hogg et al., 2013). These published data do present limitations, as they do not adequately bracket each stratigraphic unit. In an effort to maximize their utility, and estimate likely intervals of time that each unit may represent, we derived depths from the published stratigraphic diagrams and calculated a general age-depth model for the sequence. While apparently quasi-continuous, with a relatively constant depositional rate, the nature of the sequence, in terms of lithology, suggests more sporadic deposition. In plotting each unit, we have included both minimum and maximum weighted mean ages as well as potential minimum and maximum ages of the units considering potential sources of error related to radiocarbon calibration and assumptions of accumulation rates (SOM Figure S4). This highlights the clear need to initiate a systematic revision of the chronologies of many archaeological sites in the region (e.g. Loftus et al., 2016; Sealy et al., 2016) to enable more robust inter-site and inter-regional comparisons.

Taxonomic abundances (number of identified specimens: NISP) of ungulates from Boomplaas Cave were derived from Faith (2013a), with Klein's (1983) specimen counts used

for the uppermost member DGL. Comparable data (minimum number of individuals: MNI) for the Boomplaas microfauna were derived from Avery (1982). To facilitate comparison with the SWP record, we conducted a detrended correspondence analysis (DCA) for both data sets, using the first axis (rescaled from 0 to 100) to broadly summarize faunal composition (Figure 3). The linear trends in both taxonomic groups closely parallel a decline in ungulate grazers (Figure 3), suggesting that the DCA axis 1 scores are related to the replacement of grasslands by more shrubby habitats (see also Faith, 2013a). Broadly parallel changes are also observed in the south coast faunal sequences from Nelson Bay Cave (Klein, 1983) and Byneskranskop 1 (Schweitzer and Wilson, 1983) (Figure 1), implying regional shifts in habitat structure.

Humans are the primary accumulator of the Boomplaas Cave macrofauna from stratum CL4 and above (Faith, 2013a; see also Faith, 2011). This allows us to explore how the environmental changes documented at SWP translate to changes in foraging efficiency. Based on models grounded in optimal foraging theory, we examine two zooarchaeological indicators of foraging efficiency. These include the relative abundance of small-bodied and presumably low-ranked prey (leporids and tortoises) and the average food utility index (FUI; Metcalfe and Jones, 1988) – a proxy for energetic returns (e.g., meat, fat, marrow) of various ungulate body parts – of large mammal (size classes 3 and 4: 84 to 900 kg) skeletal elements (data from Faith, 2011b). Given the evidence for attrition at Boomplaas Cave (Faith and Thompson, in press), we follow Cleghorn and Marean's (2004) taphonomic model of bone survivorship and restrict analysis to the long-bones (femur, tibia, humerus, radius, and metapodials) and skull elements (crania and mandibles). Building on previous studies conducted elsewhere (e.g., Grayson, 1991, 2005; Broughton, 1994, 1999; Cannon, 2003; Faith, 2007; Munro 2004; ), we predict that declining foraging efficiency will be characterized by (1) increasing frequencies of low-ranked prey, reflecting declining availability of high-ranked prey (e.g., large game) on the landscape, and (2) increasing mean FUI values, reflecting intensified field processing due to an increase in carcass transport distances and/or search times. We recognize that because tortoises are slow-moving and easily captured, they might be considered a high-ranked prey type that should be collected whenever encountered (e.g. Stiner et al., 2000). However, the significant positive correlation between their abundances and those of leporids (Spearman's  $\rho$ :  $r_s = 0.814$ ,  $p \leq 0.001$ ), which

are unequivocally a low-ranked prey type, suggests that we can treat tortoises as low-ranked prey in this context.

<insert Table 1>

All lithic data are taken from Deacon (1982), standardized to site mean (standardized value = (layer value – site mean) / site standard deviation). We focus on the abundance of bladelets (Figure 3d), and three indicators of technological delivery: ratio of cores to retouched flakes (Figure 3g), total proportions of retouched flakes (Figure 3e), and artefact density measured as total number of artefacts per bucket of excavated sediment following Deacon (1982) (Figure 3h). These last three values function, respectively, as proxies for the transport of retouched flakes vs tool-making potential (mode of technological ‘provisioning’ (Kuhn, 1995)), frequency of flake curation, and intensity of site-use (Barton and Riel-Salvatore, 2014; Kuhn and Clark, 2015; Parry and Kelly, 1987; Riel-Salvatore and Barton, 2004). Provisioning, which we define as the systems by which stone artefact technologies are delivered in anticipation of future needs, has been argued to mediate the response of mobility to environmental change (Mackay et al., 2014; Wilkins et al., 2017). Under conditions of diminished residential mobility, we expect increases in core transport and artefact density, and lower rates of curation. The inverse is expected when mobility increases and the scheduling of movements becomes harder to predict. If bladelets are a response to diminishing subsistence returns, then their abundance should track humidity and resource productivity inversely. Similarly, if bladelets are positively associated with increasing residential mobility and declining durations of site occupancy then we expect an inverse relationship with artefact density. We also consider the relative abundance of spatially-rare, fine-grained rocks such as silcrete and crypto-crystalline silicates (CCS, subsuming chert and chalcedony) (Figure 3f); it has recently been shown that the abundance of rocks such as silcrete is responsive to increases in overall artefact abundance (Will and Mackay, 2016), and may thus reflect diminished residential mobility and improved scheduling of movements.

## Results

### *Climate change since the Last Glacial Maximum at Seweweekspoort*

The Seweweekspoort record shows substantial changes in both  $\delta^{13}\text{C}$  (range 5.1‰) and  $\delta^{15}\text{N}$  (range 9.5‰) over the last 22,300 years, implying significant changes in vegetation and climate (Figure 2). These changes are coherent with the pollen data from the same material. Across this period, a strong first-order trend is apparent, with cool, humid glacial conditions (indicated by increased cryophilic Fynbos Biome vegetation pollen and lower  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) transitioning into warmer, but substantially drier conditions during the Holocene (declining fynbos pollen and higher  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ; Figure 2). This aridification coincides with the deglacial decline in Antarctic sea-ice extent (using sea-salt sodium as a proxy) between 19-11 ka (Fischer et al., 2007; Levine et al., 2014; Wolff et al., 2010) (Figure 2), which is considered to be a strong determinant on the position of the westerly storm track (Bard and Rickaby, 2009; Stuut et al., 2004). The elevated grass pollen percentages during the last glacial period, coupled with low  $\delta^{13}\text{C}$  suggest the increased prevalence of cool growing season  $\text{C}_3$  grasses (Vogel, 1978), consistent with the model that much of the precipitation during this period fell during the winter months as a result of increased westerly influence. While recent work has indicated that variability in summer rainfall may have had some significant impact in the winter rainfall zone (Chase et al., 2015a; Chase et al., 2015b), long-term (i.e., glacial-interglacial timescales) precipitation trends in the summer rainfall zone (Chevalier and Chase, 2015) exhibit a clear antiphase relationship with humidity at Seweweekspoort (Figure 2), indicating that tropical systems played a limited role in the region at these timescales.

<insert Figure 2>

Within this broad first-order trend of deglacial aridification at Seweweekspoort, significant second-order abrupt episodes of wetter conditions (centred at 14.5, 11 and 4 cal kBP, and the last millennium) indicate major reorganisations of regional climate dynamics. In southern Africa, where rainfall regimes are defined by their strong seasonality, the varying contribution of the non-dominant moisture-bearing system can have a substantial impact on regional environments, shortening or attenuating the impact of often pronounced drought seasons (Chase et al., 2015a). In this context, humid episodes within the last glacial-interglacial transition (LGIT; 18.5-11.7 ka) – previously identified as being a period of exceptionally high effective precipitation in the region (Scholtz, 1986) – can be linked to the

warming of both high southern latitudes (Stocker, 1998; Stocker and Johnsen, 2003) and the oceans surrounding southwestern Africa (Barker et al., 2009; Farmer et al., 2005; Kim and Schneider, 2003), including a response to the slow-down of Atlantic Meridional Overturning Circulation (AMOC) during Heinrich stadial 1 (HS1; ~18-14.6 ka) (McManus et al., 2004) (Figure 2).

While the influence of the westerly storm track may have diminished as the Subtropical Front shifted poleward (Barker et al., 2009), increased evaporation from warmer oceans and the invigoration of the southern African monsoon system would have augmented the summer rain component in what was then primarily a winter rainfall regime, reducing rainfall seasonality and drought stress. At Seweweekspoort, peaks in humidity at 14.5 and 11 cal kBP typify this, with reductions in fynbos vegetation under slightly warmer conditions, and with increased grass cover as a function of more regular rains promoting shallow rooting vegetation. With the onset of the Holocene, as warming continued, the combination of tropical and temperate systems that resulted in these phases of LGIT humidity broke down. Changes in global boundary conditions resulted in 1) a more permanent southerly position of the westerly storm track, and less winter rain, and 2) strong regional warming that intensified potential evapotranspiration, enhancing drought stress (Chevalier and Chase, 2016). Combined, these factors are interpreted to have driven the marked aridification exhibited in the Seweweekspoort records (Figure 2).

### *Changing resources and technology*

The climatic changes robustly identified at Seweweekspoort are strongly reflected in our newly synthesised faunal and archaeological records (Figure 3). Large mammals and microfauna from Boomplaas Cave indicate open and grassy environments during the LGM, giving way to shrublands across the Pleistocene-Holocene transition; a phase marked by large mammal extinctions and shifts in faunal community composition throughout the region (Faith and Behrensmeyer, 2013). These changes are evident in the DCA axis 1 scores (Figure 3). The abundance of ungulate grazers and axis 1 scores at the site closely tracks  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  (Figure 3), and changes observed in the pollen record at Seweweekspoort (Figure 2). Likewise, the carbon isotope composition of tooth enamel from Boomplaas grazers parallels the  $\delta^{13}\text{C}$  shifts at Seweweekspoort, with predominantly  $\text{C}_3$  grasses consumed during the LGM giving way to increased  $\text{C}_4$  grasses during the LGIT (Sealy et al., 2016).

<insert Figure 3>

These findings contrast with earlier interpretations from floral (Deacon et al., 1984; Scholtz, 1986) and faunal assemblages (Avery, 1982; Klein, 1972; Klein, 1980, 1983) from southern Cape archaeological sequences, wherein open grassland environments – inferred primarily from a predominance of grazers in large mammal fossil records – were interpreted as signs of increased aridity. Our results support inferences that evidence for grassier vegetation indicates the influence of more/more regular precipitation on the richer soils of the valleys and plains of the southern Cape (Chase, 2010; Faith, 2013b), which currently support relatively xeric karroid vegetation (Cowling, 1983). In contemporary African ecosystems, elevated precipitation is typically associated with increased biomass of large herbivores (Coe et al., 1976; East, 1984), a phenomenon likely to have been enhanced by increased plant nutrient content under reduced atmospheric CO<sub>2</sub> concentrations (Faith, 2011a). The implication for human populations is that cooler, more humid late Pleistocene conditions presented a more productive resource base, including the proliferation of large grazing ungulates.

Zooarchaeological evidence from the human-accumulated vertebrate assemblages at Boomplaas Cave (CL4 and above) indicates that aridification through the LGIT is associated with a decline in foraging efficiency. Across the LGIT (CL4 to BRL5), tortoises and leporids increase in abundance relative to ungulates (tortoises:  $\chi^2_{\text{trend}} = 90.332$ ,  $p < 0.001$ ; leporids:  $\chi^2_{\text{trend}} = 64.854$ ,  $p < 0.001$ ). Because ungulates are larger and provide greater energetic returns than tortoises or hares – in which case they should be pursued whenever encountered – these trends imply a decline in their abundances on the landscape. This is also reflected in changes in carcass transport strategies. The average food utility index of large mammal high-survival skeletal elements increases steadily across the LGIT (Spearman's rho = 0.964,  $p < 0.001$ ), indicating a greater emphasis on skeletal parts providing the highest energetic returns. Within an optimal foraging framework, this trend is consistent with an increase in carcass transport distances and/or search times stemming from diminished availability of large game (Cannon, 2003; Faith, 2007). These parallel trends imply that the cooler and more humid conditions of the late Pleistocene provided a more productive vertebrate resource base with higher average energetic returns than did the more arid conditions of the early Holocene.

The decline in foraging efficiency through the deglaciation reverses through the Holocene. Barring an increase observed in the two upper units (Table 1), leporids decline in abundance relative to ungulates after the peak in BRL5 (~11-12 cal kBP) ( $\chi^2_{\text{trend}} = 15.854$ ,  $p < 0.001$ ). This trend, which is consistent with an increase in ungulate abundances on the landscape, complements a decline in the mean FUI of large mammal skeletal parts (Spearman's  $\rho = 0.821$ ,  $p = 0.034$ ).

From the LGM through the Holocene, there are marked temporal trends in lithic indicators of technological systems at Boomplaas Cave (Figure 3). Though not without some variation, there is a general decline through time in the abundance of bladelets ( $\chi^2_{\text{trend}} = 1323.693$ ,  $p < 0.001$ ), cores relative to retouched flakes ( $\chi^2_{\text{trend}} = 921.328$ ,  $p < 0.001$ ), and fine-grained raw materials ( $\chi^2_{\text{trend}} = 775.322$ ,  $p < 0.001$ ), coupled with an increase in the frequency of retouched pieces ( $\chi^2_{\text{trend}} = 1510.282$ ,  $p < 0.001$ ). A relationship between technological systems and the environment is suggested by strong correlations between these indicators and the DCA axis 1 scores derived for both ungulates and microfauna (Figure 3b, c; Table 2). Indeed, all the measures of technological change we employ for the Boomplaas assemblage are significantly correlated with changes in ungulate community composition, and most with the composition of micromammal communities (Table 2). Both sets of axis 1 scores track changing frequencies of grassland indicators at the site, as well as the  $\delta^{15}\text{N}$  and especially  $\delta^{13}\text{C}$  values from Seweweekspoort (Figure 3). It follows that technological change is tracking environmental change, including the patterns of climate-driven environmental change documented in the Seweweekspoort records.

<insert Table 2>

During the LGM and early LGIT bladelets are abundant and artefact indicators are consistent with low residential mobility (Figure 3). With the exception of the earliest LGM members (GWA/HCA), for which taphonomic data suggest limited human occupation (Faith, 2013a), artefact densities are well above the overall mean at Boomplaas, flake curation is uncommon, and all core to retouched flake values pre-13 ka (CL member units) are higher than all those that follow. Associated high frequencies of rock such as silcrete and CCS before 12 cal kBP (CL + BRL 7) imply regular and predictable access to fine-grained rock



through this period (Figure 3). These factors combined imply an emphasis on the transport of tool-making potential to sites.

Coincident with the onset of the first-order aridification trend after ~14 ka (BRL member units), artefact densities decrease, as do proportions of fine-grained rock and bladelets. The period is broadly characterised by more common acquisition and reduction of readily available local rocks (quartz and quartzite), with diminishing intensity of site use and little transport of cores or curation of flakes. Investment in technological costs in the later LGIT appears minimal and may reflect greater allocation of energy to search and handling of subsistence packages in response to diminishing ungulate abundance (cf., Hames, 1992; Mackay and Marwick, 2011). From the beginning of the Holocene (BRL3 through to DGL member), and tracking diminishing humidity, artefact densities are low and locally abundant rock continues to dominate, but flake curation becomes markedly more common – technological systems show a much greater emphasis on transportation and maintenance of implements in this period than in the Pleistocene.

## Conclusions

The findings presented here overturn prevailing models of environmental and behavioral change in Africa's southern Cape. A continuous and high-resolution environmental base-line is provided for the first time, indicating a trend from relative humidity during the LGM to increased aridity during the Holocene, with marked shifts in moisture across the LGIT. Rather than being characterised by 'harsh,' conditions (Deacon and Lancaster, 1988; Scholtz, 1986), the mesic environments of the late Pleistocene were highly productive, with more extensive grasslands existing in areas now dominated by drought resistant succulent shrublands. Zooarchaeological data indicate proliferation of a diverse ungulate grassland community during this time, suggesting greater resource availability for humans living in the area, and reduced search and handling times for large game. While lithic technologies track these changes, we found no evidence to support an association between the production of bladelets during the LGM/early LGIT (Robberg) and diminished subsistence conditions. Indeed, bladelets seem to have flourished in a period of relative resource abundance. The period of lowest subsistence productivity inferred from the Seweweekspoort data probably occurred during the Holocene, associated with aridification and the concomitant loss of large ungulates and faunal diversity recorded at Boomplaas. This change led to increases in carcass processing at kill sites, as evidenced by more selective transport of high utility body parts,

479 increased reliance on low-ranked prey, and a technological response in which flakes from  
480 locally acquired rocks were curated, core transportation was relatively rare, and bladelets  
481 were uncommon. In documenting the strong coupling of environmental, subsistence and  
482 technological behavior in Later Stone Age foragers, our data reflect the simple observation  
483 that all lithic technologies can be adaptive solutions, not only those often assumed to provide  
484 particular adaptive benefits. The findings afforded by high resolution analysis of late  
485 Pleistocene and Holocene climate imply more generally that the lack of certain kinds of  
486 technologies – such as bladelets, backed artefacts and bifacial points - in the earlier stages of  
487 human evolution need not carry inherent meaning. The ability of foragers to track rapid  
488 climatic and environmental changes with adaptive cultural responses is a better arbiter of  
489 cognitive complexity than the deployment of any specific technological system.

## 490 Acknowledgements

491 Funding was received from the European Research Council (ERC) under the European  
492 Union's Seventh Framework Programme (FP7/2007-2013)/ERC Starting Grant “HYRAX”,  
493 grant agreement no. 258657, and the International Union for Quaternary Research (INQUA),  
494 project no. 1205P. We would like to thank Matthew Britton and Ian Newton for assistance in  
495 collecting and analysing the material.

## 496 References

- 497 Ambrose, S.H., 2002. Small things remembered: origins of early Microlithic industries in  
498 Sub-Saharan Africa, in: Elston, R.G., Kuhn, S.L. (Eds.), *Thinking Small: Global Perspectives*  
499 *on Microlithization*. American Anthropological Association, pp. 9-30.
- 500 Ambrose, S.H., Lorenz, K.G., 1990. Social and ecological models for the Middle Stone Age  
501 in Southern Africa, in: Mellars, P. (Ed.), *The Emergence of Modern Humans*. Edinburgh  
502 University Press, Edinburgh, pp. 3-33.
- 503 Avery, D.M., 1982. Micromammals as palaeoenvironmental indicators and an interpretation  
504 of the late Quaternary in the southern Cape Province, South Africa. *Ann. S. Afr. Mus.* 85,  
505 183-377.
- 506 Bamforth, D.B., Bleed, P., 1997. Technology, flaked stone technology, and risk, in: Clark,  
507 G.A. (Ed.), *Rediscovering Darwin: Evolutionary Theory in Archaeology*. American  
508 Anthropological Association, Washington, pp. 109-140.
- 509 Bar-Yosef, O., Kuhn, S.L., 1999. The big deal about blades: Laminar technologies and  
510 human evolution. *Amer. Anthropol.* 101, 322-338.
- 511 Bard, E., Rickaby, R.E.M., 2009. Migration of the subtropical front as a modulator of glacial  
512 climate. *Nature* 460, 380-383.
- 513 Barker, S., Diz, P., Vautravers, M.J., Pike, J., Knorr, G., Hall, I.R., Broecker, W.S., 2009.  
514 Interhemispheric Atlantic seesaw response during the last deglaciation. *Nature* 457, 1097-  
515 1102.
- 516 Barton, C.M., Riel-Salvatore, J., 2014. The formation of lithic assemblages. *J. Archaeol. Sci.*  
517 46, 334-352.
- 518 Blaauw, M., Christen, J.A., 2011. Flexible Paleoclimate Age-Depth Models Using an  
519 Autoregressive Gamma Process. *Bayesian Analysis* 6, 457-474.
- 520 Bousman, C.B., 2005. Coping with risk: Later stone age technological strategies at  
521 Blydefontein Rock Shelter, South Africa. *J. Anthropol. Archaeol.* 24, 193-226.
- 522 Brown, K.S., Marean, C.W., Jacobs, Z., Schoville, B.J., Oestmo, S., Fisher, E.C., Bernatchez,  
523 J., Karkanas, P., Matthews, T., 2012. An early and enduring advanced technology originating  
524 71,000 years ago in South Africa. *Nature* 491, 590-593.
- 525 Cannon, M.D., 2003. A model of central place forager prey choice and an application to  
526 faunal remains from the Mimbres Valley, New Mexico. *J. Anthropol. Archaeol.* 22, 1-25.

527 Carr, A.S., Chase, B.M., Boom, A., Medina-Sanchez, J., 2016a. Stable isotope analyses of  
528 rock hyrax faecal pellets, hyraceum and associated vegetation in southern Africa:  
529 Implications for dietary ecology and palaeoenvironmental reconstructions. *J. Arid Environ.*  
530 134, 33-48.

531 Carr, A.S., Chase, B.M., Mackay, A., 2016b. Mid to Late Quaternary Landscape and  
532 Environmental Dynamics in the Middle Stone Age of Southern South Africa, in: Stewart,  
533 B.A., Jones, S. (Eds.), *Africa from Stages 6-2: Population Dynamics and Palaeoenvironments*.  
534 Springer.

535 Chase, B.M., 2010. South African palaeoenvironments during marine oxygen isotope stage 4:  
536 a context for the Howiesons Poort and Still Bay industries. *J. Archaeol. Sci.* 37, 1359-1366.

537 Chase, B.M., Boom, A., Carr, A.S., Carré, M., Chevalier, M., Meadows, M.E., Pedro, J.B.,  
538 Stager, J.C., Reimer, P.J., 2015a. Evolving southwest African response to abrupt deglacial  
539 North Atlantic climate change events. *QSRv* 121, 132-136.

540 Chase, B.M., Boom, A., Carr, A.S., Meadows, M.E., Reimer, P.J., 2013. Holocene climate  
541 change in southernmost South Africa: rock hyrax middens record shifts in the southern  
542 westerlies. *QSRv* 82, 199-205.

543 Chase, B.M., Lim, S., Chevalier, M., Boom, A., Carr, A.S., Meadows, M.E., Reimer, P.J.,  
544 2015b. Influence of tropical easterlies in southern Africa's winter rainfall zone during the  
545 Holocene. *QSRv* 107, 138-148.

546 Chase, B.M., Meadows, M.E., 2007. Late Quaternary dynamics of southern Africa's winter  
547 rainfall zone. *Earth-Sci. Rev.* 84, 103-138.

548 Chase, B.M., Meadows, M.E., Carr, A.S., Reimer, P.J., 2010. Evidence for progressive  
549 Holocene aridification in southern Africa recorded in Namibian hyrax middens: implications  
550 for African Monsoon dynamics and the "African Humid Period". *QuRes* 74, 36-45.

551 Chase, B.M., Meadows, M.E., Scott, L., Thomas, D.S.G., Marais, E., Sealy, J., Reimer, P.J.,  
552 2009. A record of rapid Holocene climate change preserved in hyrax middens from  
553 southwestern Africa. *Geo* 37, 703-706.

554 Chase, B.M., Quick, L.J., Meadows, M.E., Scott, L., Thomas, D.S.G., Reimer, P.J., 2011.  
555 Late glacial interhemispheric climate dynamics revealed in South African hyrax middens.  
556 *Geo* 39, 19-22.

557 Chase, B.M., Scott, L., Meadows, M.E., Gil-Romera, G., Boom, A., Carr, A.S., Reimer, P.J.,  
558 Truc, L., Valsecchi, V., Quick, L.J., 2012. Rock hyrax middens: a palaeoenvironmental  
559 archive for southern African drylands. *QSRv* 56, 107-125.

560 Chevalier, M., Chase, B.M., 2015. Southeast African records reveal a coherent shift from  
561 high- to low-latitude forcing mechanisms along the east African margin across last glacial–  
562 interglacial transition. *QSRv* 125, 117-130.

563 Chevalier, M., Chase, B.M., 2016. Determining the drivers of long-term aridity variability: a  
564 southern African case study. *JQS* 31, 143-151.

565 Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica,  
566 J.X., Hostetler, S.W., McCabe, A.M., 2009. The Last Glacial Maximum. *Sci* 325, 710-714.

567 Clarkson, C., Petraglia, M., Korisettar, R., Haslam, M., Boivin, N., Crowther, A., Ditchfield,  
568 P., Fuller, D., Miracle, P., Harris, C., Connell, K., James, H., Koshy, J., 2009. The oldest and

569 longest enduring microlithic sequence in India: 35 000 years of modern human occupation  
570 and change at the Jwalapuram Locality 9 rockshelter. *Antiquity* 83, 326–348.

571 Cleghorn, N., Marean, C.W., 2004. Distinguishing selective transport and in situ attrition: a  
572 critical review of analytical approaches. *J Taphon* 2, 43-67.

573 Coe, M.J., Cumming, D.H., Phillipson, J., 1976. Biomass and production of large African  
574 herbivores in relation to rainfall and primary production. *Oecologia* 22, 341-354.

575 Cowling, R., 1983. Phytochorology and vegetation history in the south-eastern Cape, South  
576 Africa. *J. Biogeogr.* 10, 393-419.

577 Deacon, H.J., 1979. Excavations at Boomplaas Cave : a sequence through the Upper  
578 Pleistocene and Holocene in South Africa. *Wld. Archaeol.* 10, 241-257.

579 Deacon, H.J., Deacon, J., Scholtz, A., Thackeray, J.F., Brink, J.S., 1984. Correlation of  
580 palaeoenvironmental data from the Late Pleistocene and Holocene deposits at Boomplaas  
581 Cave, southern Cape, in: Vogel, J.C. (Ed.), *Late Cainozoic Palaeoclimates of the Southern*  
582 *Hemisphere*. Balkema, Rotterdam, pp. 339-360.

583 Deacon, J., 1972. Wilton: An assessment after fifty years. *S. Afr. Archaeol. Bull.* 27, 10-48.

584 Deacon, J., 1978. Changing patterns in late Pleistocene / early Holocene prehistory in  
585 southern Africa as seen from the Nelson Bay Cave stone artifact sequence. *QuRes* 10, 84-  
586 111.

587 Deacon, J., 1982. *The Later Stone Age in the southern Cape, South Africa*. University of  
588 Cape Town, Cape Town, South Africa.

589 Deacon, J., 1984. *The Later Stone Age in southernmost Africa*. Archaeopress, Oxford.

590 Deacon, J., Lancaster, N., 1988. *Late Quaternary palaeoenvironments of southern Africa*.  
591 Clarendon Press, Oxford.

592 Delagnes, A., Schmidt, P., Douze, K., Wurz, S., Bellot-Gurlet, L., Conard, N.J., Nickel, K.G.,  
593 van Niekerk, K.L., Henshilwood, C.S., 2016. Early Evidence for the Extensive Heat  
594 Treatment of Silcrete in the Howiesons Poort at Klipdrift Shelter (Layer PBD, 65 ka), South  
595 Africa. *PLoS One* 11, e0163874.

596 East, R., 1984. Rainfall, soil nutrient status and biomass of large African savanna mammals.  
597 *Afr. J. Ecol.* 22, 245-270.

598 Ehleringer, J.R., Cooper, T.A., 1988. Correlations between carbon isotope ratio and  
599 microhabitat of desert plants. *Oecologia* 76, 562-566.

600 Elston, R.G.-., Brantingham, P.J., 2002. Microlithic technology in Northern Asia: A risk-  
601 minimizing strategy of the Late Paleolithic and Early Holocene, in: Elston, R.G., Kuhn, S.L.  
602 (Eds.), *Thinking Small: Global Perspectives on Microlithization*.

603 Faith, J.T., 2007. Changes in reindeer body part representation at Grotte XVI, Dordogne,  
604 France. *J. Archaeol. Sci.* 34, 2003-2011.

605 Faith, J.T., 2011a. Late Pleistocene climate change, nutrient cycling, and the megafaunal  
606 extinctions in North America. *QSRv* 30, 1675-1680.

607 Faith, J.T., 2011b. *Late Quaternary Megafaunal Extinctions in Southern Africa's Cape Floral*  
608 *Region*, Department of Anthropology. The George Washington University.

609 Faith, J.T., 2013a. Taphonomic and paleoecological change in the large mammal sequence  
610 from Boomplaas Cave, Western Cape, South Africa. *J. Hum. Evol.* 65, 715-730.

611 Faith, J.T., 2013b. Ungulate diversity and precipitation history since the Last Glacial  
612 Maximum in the Western Cape, South Africa. *QSRv* 68, 191-199.

613 Faith, J.T., Behrensmeyer, A.K., 2013. Climate change and faunal turnover: testing the  
614 mechanics of the turnover-pulse hypothesis with South African fossil data. *Palb* 39, 609-627.

615 Faith, J.T., Thompson, J.C., in press. Low-survival skeletal elements track attrition, not  
616 carcass transport behavior in Quaternary large mammal assemblages, in: Giovas, C.M.,  
617 LeFebvre, M. (Eds.), *Zooarchaeology in Practice: Case Studies in Methodology and*  
618 *Interpretation in Archaeofaunal Analysis*. Springer.

619 Farmer, E.C., deMenocal, P.B., Marchitto, T.M., 2005. Holocene and deglacial ocean  
620 temperature variability in the Benguela upwelling region: implications for low-latitude  
621 atmospheric circulation. *PalOc* 20, doi:10.1029/2004PA001049.

622 Farquhar, G.D., Ehleringer, J.R., Hubick, K.T., 1989. Carbon isotope discrimination and  
623 photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 40, 503-537.

624 Farquhar, G.D., Richards, R.A., 1984. Isotopic composition of plant carbon correlates with  
625 water-use efficiency of wheat genotypes. *Aust. J. Plant Physiol.* 11, 539-552.

626 Fischer, H., Fundel, F., Ruth, U., Twarloh, B., Wegner, A., Udisti, R., Becagli, S., Castellano,  
627 E., Morganti, A., Severi, M., Wolff, E., Littot, G., Röthlisberger, R., Mulvaney, R., Hutterli,  
628 M.A., Kaufmann, P., Federer, U., Lambert, F., Bigler, M., Hansson, M., Jonsell, U., de  
629 Angelis, M., Boutron, C., Siggaard-Andersen, M.-L., Steffensen, J.P., Barbante, C., Gaspari,  
630 V., Gabrielli, P., Wagenbach, D., 2007. Reconstruction of millennial changes in dust  
631 emission, transport and regional sea ice coverage using the deep EPICA ice cores from the  
632 Atlantic and Indian Ocean sector of Antarctica. *E&PSL* 260, 340-354.

633 Foley, R.A., Lahr, M.M., 2003. On stony ground: Lithic technology, human evolution, and  
634 the emergence of culture. *Evol Anthropol* 12, 109-122.

635 Goebel, T., 2002. The "Microblade Adaptation" and recolonization of Siberia during the Late  
636 Upper Pleistocene, in: Elston, R.G., Kuhn, S.L. (Eds.), *Thinking Small: Global Perspectives*  
637 *on Microlithization*.

638 Grosjean, M., Cartajena, I., Geyh, M.A., Nunez, L., 2003. From proxy data to paleoclimate  
639 interpretation: the mid-Holocene paradox of the Atacama Desert, northern Chile.  
640 *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 194, 247-258.

641 Hames, R.B., 1992. Time allocation, in: Smith, E.A., Winterhalder, B. (Eds.), *Evolutionary*  
642 *Ecology and Human Behavior*. Aldine de Gruyter, New York, pp. 203-235.

643 Handley, L.L., Austin, A.T., Stewart, G.R., Robinson, D., Scrimgeour, C.M., Raven, J.A.,  
644 Heaton, T.H.E., Schmidt, S., 1999. The  $\delta^{15}\text{N}$  natural abundance ( $\delta^{15}\text{N}$ ) of ecosystem samples  
645 reflects measures of water availability. *Funct. Plant Biol.* 26, 185-199.

646 Handley, L.L., Odee, D., Scrimgeour, C.M., 1994.  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  patterns in savanna  
647 vegetation: dependence on water availability and disturbance. *Funct. Ecol.* 8, 306-314.

648 Hartman, G., 2011. Are elevated  $\delta^{15}\text{N}$  values in herbivores in hot and arid environments  
649 caused by diet or animal physiology? *Funct. Ecol.* 25, 122-131.

650 Heaton, T.H.E., 1987. The  $^{15}\text{N}/^{14}\text{N}$  ratios of plants in South Africa and Namibia: relationship  
651 to climate and coastal/saline environments. *Oecologia* 74, 236-246.

652 Heaton, T.H.E., Talma, A.S., Vogel, J.C., 1986. Dissolved gas paleotemperatures and  $^{18}\text{O}$   
653 variations derived from groundwater near Uitenhage, South Africa. *QuRes* 25, 79-88.

654 Henshilwood, C., d'Errico, F., Vanhaeren, M., Van Niekerk, K., Jacobs, Z., 2004a. Middle  
655 Stone Age shell beads from South Africa. *Sci* 304, 404.

656 Henshilwood, C.S., d'Errico, F., Vanhaeren, M., van Niekerk, K., Jacobs, Z., 2004b. Middle  
657 Stone Age shell beads from South Africa. *Sci* 384, 404.

658 Henshilwood, C.S., d'Errico, F., Yates, R., Jacobs, Z., Tribolo, C., Duller, G.A.T., Mercier,  
659 N., Sealy, J.C., Valladas, H., Watts, I., Wintle, A.G., 2002. Emergence of modern human  
660 behavior: Middle Stone Age engravings from South Africa. *Sci* 295, 1278-1280.

661 Henshilwood, C.S., van Niekerk, K.L., Wurz, S., Delagnes, A., Armitage, S.J., Rifkin, R.F.,  
662 Douze, K., Keene, P., Haaland, M.M., Reynard, J., Discamps, E., Mienies, S.S., 2014.  
663 Klipdrift Shelter, southern Cape, South Africa: Preliminary Report on the Howiesons Poort  
664 layers. *J. Archaeol. Sci.* 45, 284-303.

665 Hogg, A.G., Hua, Q., Blackwell, P.G., Niu, M., Buck, C.E., Guilderson, T.P., Heaton, T.J.,  
666 Palmer, J.G., Reimer, P.J., Reimer, R.W., Turney, C.S.M., Zimmerman, S.R.H., 2013.  
667 SHCal13 Southern Hemisphere Calibration, 0–50,000 Years cal BP.

668 Kim, J.-H., Schneider, R.R., 2003. Low-latitude control of interhemispheric sea-surface  
669 temperature contrast in the tropical Atlantic over the past 21 kyrs: the possible role of SE  
670 trade winds. *CIDy* 23, 337-347.

671 Klein, R.G., 1972. The late Quaternary mammalian fauna of Nelson Bay Cave (Cape  
672 Province, South Africa): its implications for megafaunal extinctions and environmental and  
673 cultural change. *QuRes* 2, 135-142.

674 Klein, R.G., 1980. Environmental and ecological implications of large mammals from Upper  
675 Pleistocene and Holocene sites in southern Africa. *Ann. S. Afr. Mus.*, 223-283.

676 Klein, R.G., 1983. Palaeoenvironmental implications of Quaternary large mammals in the  
677 Fynbos region, in: Deacon, H.J., Hendey, Q.B. and Lambrechts, J.J.N. (Ed.), *Fynbos*  
678 *Palaeoecology: a preliminary synthesis*, 75 ed, pp. 116-138.

679 Kohfeld, K.E., Graham, R.M., de Boer, A.M., Sime, L.C., Wolff, E.W., Le Quéré, C., Bopp,  
680 L., 2013. Southern Hemisphere westerly wind changes during the Last Glacial Maximum:  
681 paleo-data synthesis. *QSRv* 68, 76-95.

682 Kuhn, S.L., 1995. *Mousterian Lithic Technology*. Princeton University Press, Princeton.

683 Kuhn, S.L., Clark, A.E., 2015. Artifact densities and assemblage formation: Evidence from  
684 Tabun Cave. *J. Anthropol. Archaeol.* 38, 8-16.

685 Levine, J.G., Yang, X., Jones, A.E., Wolff, E.W., 2014. Sea salt as an ice core proxy for past  
686 sea ice extent: a process-based model study. *Journal of Geophysical Research: Atmospheres*  
687 119, 5737-5756.

688 Loftus, E., Sealy, J., Lee-Thorp, J., 2016. New Radiocarbon Dates and Bayesian Models for  
689 Nelson Bay Cave and Byneskranskop 1: Implications for the South African Later Stone Age  
690 Sequence. *Radiocarbon FirstView*, 1-17.

691 Lombard, M., Wadley, L., Deacon, J., Wurz, S., Parsons, I., Mohapi, M., Swart, J., Mitchell,  
692 P., 2012. South African and Lesotho Stone Age sequence updated. *S. Afr. Archaeol. Bull.* 67,  
693 120-144.

694 Lowe, J.J., Rasmussen, S.O., Björck, S., Hoek, W.Z., Steffensen, J.P., Walker, M.J.C., Yu,  
695 Z.C., 2008. Synchronisation of palaeoenvironmental events in the North Atlantic region  
696 during the Last Termination: a revised protocol recommended by the INTIMATE group.  
697 *QSRv* 27, 6-17.

698 Mackay, A., 2009. History and Selection in the Late Pleistocene Archaeology of the Western  
699 Cape, South Africa, School of Archaeology and Anthropology. Australian National  
700 University.

701 Mackay, A., Marwick, B., 2011. Costs and benefits of technological decision making under  
702 variable conditions: Examples from the late Pleistocene in southern Africa, in: Marwick, B.,  
703 Mackay, A. (Eds.), *Keeping your Edge: Recent Approaches to the Organisation of Stone*  
704 *Artefact Technology*. Archaeopress, Oxford, pp. 119-134.

705 Mackay, A., Stewart, B.A., Chase, B.M., 2014. Coalescence and fragmentation in the late  
706 Pleistocene archaeology of southernmost Africa. *J. Hum. Evol.* 72, 26-51.

707 Marean, C.W., 2010. Pinnacle Point Cave 13B (Western Cape Province, South Africa) in  
708 context: The Cape Floral kingdom, shellfish, and modern human origins. *J. Hum. Evol.* 59,  
709 425-443.

710 Marean, C.W., 2014. The origins and significance of coastal resource use in Africa and  
711 Western Eurasia. *J. Hum. Evol.* 77, 17-40.

712 Marean, C.W., Cawthra, H.C., Cowling, R.M., Esler, K.J., Fisher, E., Milewski, A., Potts,  
713 A.J., Singels, E., De Vynck, J., 2014. Stone Age people in a changing South African Greater  
714 Cape Floristic Region, in: N. Allsopp, Colville, J.F., Verboom, G.A. (Eds.), *Fynbos: Ecology,*  
715 *Evolution, and Conservation of a Megadiverse Region*. Oxford University Press, Oxford, UK,  
716 pp. 164-199.

717 McCall, G.S., 2007. Behavioral ecological models of lithic technological change during the  
718 later Middle Stone Age of South Africa. *J. Archaeol. Sci.* 34, 1738-1751.

719 McCall, G.S., Thomas, J.T., 2012. Still Bay and Howiesons Poort Foraging Strategies:  
720 Recent Research and Models of Culture Change. *Afr. Archaeol. Rev.* 29, 7-50.

721 McManus, J.F., Francois, R., Gherardi, J.-M., Keigwin, L.D., Brown-Leger, S., 2004.  
722 Collapse and rapid resumption of Atlantic meridional circulation linked to deglacial climate  
723 changes. *Nature* 428, 834-837.

724 Mellars, P., 2006. Why did modern human populations disperse from Africa ca. 60,000 years  
725 ago? A new model. *Proc. Natl. Acad. Sci. U.S.A.* 103, 9381-9386.

726 Metcalfe, D., Jones, K.T., 1988. A Reconsideration of Animal Body-Part Utility Indices.  
727 *Amer. Antiq.* 53, 486-504.

728 Mitchell, P., 2000. The organisation of Later Stone Age lithic technology in the Caledon  
729 Valley, Southern Africa. *Afr. Archaeol. Rev.* 17, 141-176.

730 Mitchell, P.J., 1988. *The early microlithic assemblages of southern Africa*. BAR, Oxford.

731 Moffett, R.O., Deacon, H.J., 1977. The Flora and Vegetation in the Surrounds of Boomplaas  
732 Cave: Cango Valley. *The South African Archaeological Bulletin* 32, 127-145.



733 Moore, P.D., Webb, J.A., Collinson, M.E., 1991. Pollen Analysis, 2nd ed. Blackwell  
734 Scientific Publications, Oxford.

735 Mucina, L., Rutherford, M.C., 2006. The vegetation of South Africa, Lesotho and Swaziland,  
736 Strelitzia. South African National Biodiversity Institute, Pretoria.

737 Murphy, B.P., Bowman, D.M.J.S., 2006. Kangaroo metabolism does not cause the  
738 relationship between bone collagen  $\delta^{15}\text{N}$  and water availability. *Funct. Ecol.* 20, 1062-1069.

739 Murphy, B.P., Bowman, D.M.J.S., 2009. The carbon and nitrogen isotope composition of  
740 Australian grasses in relation to climate. *Funct. Ecol.* 23, 1040-1049.

741 Neeley, M.P., 2002. Going microlithic: A Levantine perspective on the adoption of  
742 microlithic technologies, in: Elston, R.G., Kuhn, S.L. (Eds.), *Thinking Small: Global*  
743 *Perspectives on Microlithization*.

744 Pargeter, J., 2016. Lithic miniaturization in Late Pleistocene southern Africa. *Journal of*  
745 *Archaeological Science: Reports* 10, 221-236.

746 Parkington, J., Cartwright, C., Cowling, R.M., Baxter, A., Meadows, M., 2000.  
747 Palaeovegetation at the Last Glacial Maximum in the Western Cape, South Africa: wood  
748 charcoal and pollen evidence from Elands Bay Cave. *SAJSc* 96, 543-546.

749 Parry, W.J., Kelly, R.L., 1987. Expedient core technology and sedentism, in: Johnson, J.K.,  
750 Morrow, C.A. (Eds.), *The organization of core technology*. Westview Press, pp. 285-304.

751 Partridge, T.C., Scott, L., Hamilton, J.E., 1999. Synthetic reconstructions of southern African  
752 environments during the Last Glacial Maximum (21-18 kyr) and the Holocene Altithermal  
753 (8-6 kyr). *Quat Int* 57-8, 207-214.

754 Partridge, T.C., Scott, L., Schneider, R.R., 2004. Between Agulhas and Benguela: responses  
755 of southern African climates of the late Pleistocene to current fluxes, orbital precession and  
756 the extent of the circum-Antarctic vortex, *Past Climate Variability through Europe and*  
757 *Africa*, pp. 45-68.

758 Pate, J.S., 2001. Carbon isotope discrimination and plant water-use efficiency: case scenarios  
759 for  $\text{C}_3$  plants, in: Unkovich, M., Pate, J., McNeill, A., Gibbs, D.J. (Eds.), *Stable Isotope*  
760 *Techniques in the Study of Biological Processes and Functioning of Ecosystems*. Kluwer  
761 Academic Publishers, Dordrecht, pp. 19-37.

762 Petraglia, M., Clarkson, C., Boivin, N., Haslam, M., Korisettar, R., Chaubey, G., Ditchfield,  
763 P., Fuller, D., James, H., Jones, S., Kivisild, T., Koshy, J., Lahr, M.M., Metspalu, M.,  
764 Roberts, R., Arnold, L., 2009. Population increase and environmental deterioration  
765 correspond with microlithic innovations in South Asia ca. 35,000 years ago. *Proc Natl Acad*  
766 *Sci U S A* 106, 12261-12266.

767 Powell, A., Shennan, S., Thomas, M.G., 2009. Late Pleistocene demography and the  
768 appearance of modern human behavior. *Sci* 324, 1298-1301.

769 Quick, L.J., Carr, A.S., Meadows, M.E., Boom, A., Bateman, M.D., Roberts, D.L., Reimer,  
770 P.J., Chase, B.M., 2015. A late Pleistocene–Holocene multi-proxy record of  
771 palaeoenvironmental change from Still Bay, southern Cape Coast, South Africa. *JQS* 30, 870-  
772 885.

773 Quick, L.J., Meadows, M.E., Bateman, M.D., Kirsten, K.L., Mäusbacher, R., Haberzettl, T.,  
774 Chase, B.M., 2016. Vegetation and climate dynamics during the last glacial period in the

775 fynbos-afrotemperate forest ecotone, southern Cape, South Africa. *Quat Int* 404, Part B, 136-  
776 149.

777 Riel-Salvatore, J., Barton, M., 2004. Late Pleistocene Technology, Economic Behavior, and  
778 Land-Use Dynamics in Southern Italy. *Amer. Antiq.* 69, 257-274.

779 Roberts, P., Henshilwood, C.S., van Niekerk, K.L., Keene, P., Gledhill, A., Reynard, J.,  
780 Badenhorst, S., Lee-Thorp, J., 2016. Climate, environment and early human innovation:  
781 stable isotope and faunal proxy evidence from archaeological Sites (98-59ka) in the Southern  
782 Cape, South Africa. *PLoS One* 11, e0157408.

783 Rutherford, M.C., Mucina, L., Powrie, L.W., 2012. The South African National Vegetation  
784 Database: History, development, applications, problems and future. *SAJSc* 108.

785 Rutherford, M.C., Powrie, L.W., Midgley, G.F., 2003. ACKDAT: A digital spatial database  
786 of distributions of South African plant species and species assemblages. *S. Afr. J. Bot.* 69,  
787 99-104.

788 Sale, J.B., 1965. The feeding behaviour of rock hyraxes (genera *Procavia* and *Heterohyrax*)  
789 in Kenya. *Afr. J. Ecol.* 3, 1-18.

790 SANBI, 2003. PRECIS (National Herbarium Pretoria (PRE) Computerized Information  
791 System) database.

792 Scholtz, A., 1986. Palynological and Palaeobotanical Studies in the Southern Cape.  
793 University of Stellenbosch, Stellenbosch, South Africa.

794 Schweitzer, F.R., Wilson, M.L., 1983. Byneskranskop 1: A late Quaternary living site in the  
795 southern Cape province, South Africa. The Rustica Press, Cape Town.

796 Scott, L., 1982. Late Quaternary fossil pollen grains from the Transvaal, South Africa. *Rev.*  
797 *Palaeobot. Palynol.* 36, 241-278.

798 Sealy, J., Lee-Thorp, J., Loftus, E., Faith, J.T., Marean, C.W., 2016. Late Quaternary  
799 environmental change in the Southern Cape, South Africa, from stable carbon and oxygen  
800 isotopes in faunal tooth enamel from Boomplaas Cave. *JQS* 31, 919-927.

801 Sime, L.C., Kohfeld, K.E., Le Quéré, C., Wolff, E.W., de Boer, A.M., Graham, R.M., Bopp,  
802 L., 2013. Southern Hemisphere westerly wind changes during the Last Glacial Maximum:  
803 model-data comparison. *QSRv* 64, 104-120.

804 Smith, B.N., 1972. Natural abundance of the stable isotopes of carbon in biological systems.  
805 *BioSc* 22, 226-231.

806 Stiner, M.C., Munro, N.D., Surovell, T.A., 2000. The tortoise and the hare: small-game use,  
807 the broad spectrum revolution, and Paleolithic demography. *Curr. Anthropol.* 41, 39-73.

808 Stocker, T.F., 1998. The seesaw effect. *Sci* 282, 61-62.

809 Stocker, T.F., Johnsen, S.J., 2003. A minimum thermodynamic model for the bipolar seesaw.  
810 *PalOc* 18.

811 Stockmarr, J., 1971. Tablets with spores used in absolute pollen analysis. *Pollen et Spores* 13,  
812 614-621.

813 Stone, A.E.C., 2014. Last Glacial Maximum conditions in southern Africa: are we any closer  
814 to understanding the climate of this time period? *Progress in Physical Geography*.

815 Stuut, J.-B.W., Crosta, X., Van der Borg, K., Schneider, R.R., 2004. On the relationship  
816 between Antarctic sea ice and southwestern African climate during the late Quaternary. *Geo*  
817 32, 909-912.

818 Talma, A.S., Vogel, J.C., 1992. Late Quaternary paleotemperatures derived from a  
819 speleothem from Cango Caves, Cape Province, South Africa. *QuRes* 37, 203-213.

820 van Zinderen Bakker, E.M., 1953. South African pollen grains and spores. Part I. Balkema,  
821 Amsterdam/Cape Town.

822 van Zinderen Bakker, E.M., 1956. South African pollen grains and spores. Part II. Balkema,  
823 Amsterdam/Cape Town.

824 van Zinderen Bakker, E.M., Coetzee, J.A., 1959. South African pollen grains and spores. Part  
825 III. Balkema, Amsterdam/Cape Town.

826 Vogel, J.C., 1978. Isotopic assessment of the dietary habits of ungulates. *SAJSc* 74, 298-301.

827 Wadley, L., 1993. The Pleistocene Later Stone Age south of the Limpopo River. *Journal of*  
828 *World Prehistory* 7, 243-296.

829 Wang, L., D'Odorico, P., Ries, L., Macko, S.A., 2010. Patterns and implications of plant-soil  
830  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in African savanna ecosystems. *QuRes* 73, 77-83.

831 Wilkins, J., Brown, K.S., Oestmo, S., Pereira, T., Ranhorn, K.L., Schoville, B.J., Marean,  
832 C.W., 2017. Lithic technological responses to Late Pleistocene glacial cycling at Pinnacle  
833 Point Site 5-6, South Africa. *PLoS One* 12, e0174051.

834 Will, M., Mackay, A., 2016. What factors govern the procurement and use of silcrete during  
835 the Stone Age of South Africa? *Journal of Archaeological Science: Reports*.

836 Wolff, E.W., Barbante, C., Becagli, S., Bigler, M., Boutron, C.F., Castellano, E., de Angelis,  
837 M., Federer, U., Fischer, H., Fundel, F., Hansson, M., Hutterli, M., Jonsell, U., Karlin, T.,  
838 Kaufmann, P., Lambert, F., Littot, G.C., Mulvaney, R., Röthlisberger, R., Ruth, U., Severi,  
839 M., Siggaard-Andersen, M.L., Sime, L.C., Steffensen, J.P., Stocker, T.F., Traversi, R.,  
840 Twarloh, B., Udisti, R., Wagenbach, D., Wegner, A., 2010. Changes in environment over the  
841 last 800,000 years from chemical analysis of the EPICA Dome C ice core. *QSRv* 29, 285-  
842 295.

843

## 844 Table Captions

845 **Table 1:** Zooarchaeological indicators of foraging efficiency at Boomplaas Cave. These include the  
846 relative abundance of tortoises (%), leporids (%), and mean food utility index (FUI) of high-survival  
847 body parts belonging to size 3-4 mammals.

848 **Table 2:** The correlation (Spearman's  $\rho$ ) between lithic technological indicators and faunal  
849 community composition (DCA Axis 1 scores) for ungulates and microfauna at Boomplaas Cave.  
850 Significant values in bold.

## 851 Figure Captions

**Figure 1:** (A) Map of southern Africa showing seasonality of rainfall and climatic gradients dictated by the zones of summer/tropical (orange) and winter/temperate (blue) rainfall dominance. Winter rainfall is primarily a result of frontal systems embedded in the westerly storm track. Major atmospheric (white arrows) and oceanic (blue arrows) circulation systems and the austral summer positions of the Inter-Tropical Convergence Zone (ITCZ) and the Congo Air Boundary (CAB) are indicated. The location of the study site in the transitional southern Cape region is shown. (B) Map of southwest African coastal region with the Seweweekspoort sites and other key palaeoenvironmental and archaeological sites indicated (shading as for panel 'A'). (C) Topographical map of Seweweekspoort, with the SWP-1 and SWP-3 sites indicated.

**Figure 2:** Comparison of  $\delta^{15}\text{N}$  (e),  $\delta^{13}\text{C}$  values (b), fynbos (c) and grass (d) pollen percentages from the Seweweekspoort hyrax middens with relevant palaeoenvironmental records including the northern summer rainfall zone wet season precipitation reconstruction (a; Chevalier and Chase, 2015), the Bermuda Rise record of Atlantic Meridional Overturning Circulation (AMOC) strength and the northward oceanic transport of heat (f; McManus et al., 2004), foraminifera records indicating conditions in the ocean to the south of the Africa (g, h; Barker et al., 2009) and sea salt sodium concentrations from the EPICA DML ice core in Antarctica (i; Fischer et al., 2007).

**Figure 3:** Comparison of  $\delta^{15}\text{N}$  (j) and  $\delta^{13}\text{C}$  (i) values from the Seweweekspoort hyrax middens with zooarchaeological records (a-c) of macrofauna (Faith, 2013a; Klein, 1983) and microfauna (Avery, 1982), and (d-h) lithic data (Deacon, 1979; Deacon et al., 1984) from Boomplaas Cave. Error bars on the data from Boomplaas reflect potential age ranges ( $2\sigma$ ) of each stratigraphic unit.